

B121

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
9 February 2006 (09.02.2006)

PCT

(10) International Publication Number
WO 2006/014519 A1(51) International Patent Classification⁷: C12Q 1/00(21) International Application Number:
PCT/US2005/024046

(22) International Filing Date: 7 July 2005 (07.07.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/586,043 7 July 2004 (07.07.2004) US(71) Applicant (for all designated States except US): THE
GENERAL HOSPITAL CORPORATION [US/US]; 55
Fruit Street, Boston, MA 02114 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): ELMALEH, David,
R. [US/US]; 38 Hartman Road, Newton, MA 02159 (US).(74) Agents: ZARUTSKIE, Jennifer, A. et al.; Patent Group,
Foley Hoag LLP, 155 Seaport Boulevard, Boston, MA
02210-2600 (US).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,
MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ,
OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL,
SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC,
VN, YU, ZA, ZM, ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: DIRECT ACTIVATION OF ATIII IN WHOLE BLOOD AND PLASMA

(57) Abstract: Methods of activating ATIII *in situ* in a blood product are disclosed, as is the use of such methods and blood products
in treating infectious diseases, inflammatory disorders and diseases or conditions that are mediated by thrombin activation.

WO 2006/014519 A1

WO 2006/014519

PCT/US2005/024046

DIRECT ACTIVATION OF ATIII IN WHOLE BLOOD AND PLASMA**RELATED APPLICATIONS**

- 5 [0001] This application claims priority to provisional application U.S.S.N. 60/586,043, filed July 7, 2004, the content of which application is specifically incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

- 10 [0002] Although vaccines are available to prevent many types of viral infection, not all viruses are able to be prevented by a vaccine and not all potential victims are able to receive vaccinations even if they are available. For example, the retrovirus human immunodeficiency virus (HIV) causes Acquired Immunodeficiency Syndrome (AIDS), an incurable disease for which there is no vaccine in which the body's immune system breaks
15 down leaving the victim vulnerable to opportunistic infections, *e.g.*, pneumonia, and certain cancers, *e.g.*, Kaposi's Sarcoma. Many patients with HIV are co-infected with Hepatitis C Virus (HCV), Hepatitis B (HBV), or other viruses. Although a vaccine is available for certain viruses such as HBV, many at-risk people do not receive or have access to the vaccine. Further, some data indicate that the HBV vaccine is not as effective in people
20 already infected with HIV.

- [0003] Viral infections, once established, are generally incurable. There are, however, a variety of anti-viral drugs that can prevent viruses from reproducing and ravaging the body's immune system, *i.e.*, that slow the infection and lengthen the subject's life. However, such therapies often only partially effective, and it is unknown how much viral
25 suppression is required to achieve durable virologic, immunologic, and clinical benefits. Anti-viral drugs are often highly toxic and can cause serious side effects, including heart damage, kidney failure, and osteoporosis.

- [0004] For example, highly active antiretroviral drug therapy (HAART) is a widely used anti-HIV therapy that entails multiple-drug protease inhibitor-containing regimens that can
30 completely suppress viral replication. Hepatic injury is a major concern as a result of antiretroviral therapy (HAART) and has been shown to occur with all classes of antiretroviral therapy. The efficacy of current anti-HIV therapy is further limited by the complexity of regimens, pill burden, and drug-drug interactions. Compliance with the toxic effects of antiretroviral drugs make a lifetime of combination therapy a difficult prospect

WO 2006/014519

PCT/US2005/024046

and many patients cannot tolerate long-term treatment with HAART. Further, poor adherence to combination therapy regimes has led to the emergence of drug-resistant strains of HIV.

5 [0005] There is clearly a need for new anti-viral agents and other novel approaches to treating viral infection.

SUMMARY OF THE INVENTION

10 [0006] The present invention is based on the surprising finding that ATIII may be activated using saccharides such as heparin directly *in situ* in a blood product, such as blood, plasma, serum albumin, recombinant plasma, and the like, and re-administered to a subject. For example, a blood product of a subject may be incubated with heparin or other saccharide, dialyzed, and re-administered into the subject. Alternatively, a blood product from another source, such as a commercial source, may be incubated with heparin or other saccharide, dialyzed, and administered to a subject. The blood products so treated contain activated

15 ATIII in an effective dose to treat viral infection. This novel treatment eliminates the need for activated ATIII preparation in advance. Such treated blood products as a treatment may be useful, for example, in treating patients having viral infections or diseases or conditions that are caused by or contributed to by thrombin activation. The methods of treatment provided by the present invention may result in shorter pre-clinical and clinical testing

20 times. Further, these methods of treatment provide an alternative, or supplement, to existing treatment methods of viral infection using purified activated ATIII pharmaceutical preparations.

[0007] Other features and advantages of the invention will become apparent from the following detailed description and claims.

25

BRIEF DESCRIPTION OF THE DRAWINGS

[0008] FIGURE 1 depicts the HIV-1 inhibitory activity in blood treated with LMW heparin (as described below in Example 4).

30 [0009] FIGURE 2 depicts the mechanism of action of activated heparin. Pure ATIII from Aventis (human source) and from GTC (recombinant) are inactive. Oligosaccharide and ATIII incubation and/or incubation with heat accelerate the interaction of these molecules to form a complex that undergoes a conformational change to produce an active form.

WO 2006/014519

PCT/US2005/024046

DETAILED DESCRIPTION OF THE INVENTION**[0010] 1. Definitions**

[0011] For convenience, certain terms employed in the specification, examples, and appended claims are collected here. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.

[0012] The articles "a" and "an" are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

[0013] The term "activating ATIII" or "to activate ATIII" refers to treating ATIII under suitable conditions such that the ATIII becomes high molecular weight ATIII and is able to reduce viral load, i.e."

[0014] The term "administering" includes any method of delivery of a pharmaceutical composition or therapeutic agent into a subject's system or to a particular region in or on a subject. The phrases "systemic administration," "administered systemically," "peripheral administration" and "administered peripherally" as used herein mean the administration of a compound, drug or other material other than directly into the central nervous system, such that it enters the patient's system and, thus, is subject to metabolism and other like processes, for example, subcutaneous administration. "Parenteral administration" and "administered parenterally" means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intra-articular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

[0015] The term "ATIII" refers to antithrombin III.

[0016] The term "blood product" refers to any product or substance that is blood or is derived from blood. For example, blood products include, but are not limited to, whole blood, plasma, serum, serum albumin preparations, and artificial preparations of the same, such as recombinant plasma.

[0017] The term "mammal" is known in the art, and exemplary mammals include humans, primates, bovines, porcines, canines, felines, and rodents (e.g., mice and rats).

WO 2006/014519

PCT/US2005/024046

[0018] The term "high molecular weight ATIII" means ATIII that has been treated to increase the molecular weight of the molecule relative to wildtype ATIII, e.g. to about 60 Kda to about 240 kDa from 58 kDa.

5 [0019] A "patient," "subject" or "host" to be treated by the subject method may mean either a human or non-human animal.

[0020] The phrase "pharmaceutically acceptable" refers to those compositions and dosages thereof within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

10 [0021] The phrase "pharmaceutically-acceptable carrier" means a pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting any supplement or composition, or component thereof, from one organ, or portion of the body, to another organ, or portion of the body. The term "pharmaceutically acceptable carrier"

15 refers to a carrier(s) that is "acceptable" in the sense of being compatible with the other ingredients of a composition and not deleterious to the recipient thereof. Some examples of materials which may serve as pharmaceutically-acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and

20 cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as

25 magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations.

[0022] The term "saccharide" includes both monosaccharides and polysaccharides.

30 [0023] The term "a saccharide able to activate ATIII" refers to any saccharide able to transform wildtype ATIII into activated and/or high molecular weight ATIII that has the ability to reduce viral load.

WO 2006/014519

PCT/US2005/024046

[0024] The term "therapeutically effective amount" refers to that amount of activated ATIII, drug or other molecule which is sufficient to effect treatment when administered to a subject in need of such treatment. The therapeutically effective amount will vary depending upon the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of ordinary skill in the art.

[0025] The term "treating" as used herein is intended to encompass curing as well as ameliorating at least one symptom or the progression of any condition or disease. decreased or prevented. Treatment may be performed either prophylactically, or subsequent to the initiation of a pathologic event.

[0026] The term "viral load" refers to the concentration of a virus, such as HIV, in the blood.

[0027] Unless otherwise indicated, all numbers expressing quantities of ingredients, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term "about." Accordingly, unless indicated to the contrary, the numerical parameters set forth in this specification and attached claims are approximations that may vary depending upon the desired properties sought to be obtained by the present invention.

[0028] 2. In Situ Blood Activation of ATIII

[0029] Antithrombin III (ATIII) is a glycoprotein present in blood plasma with a well-defined role in blood clotting. Specifically, ATIII is a potent inhibitor of the reactions of the coagulation cascade with an apparent molecular weight of between 54k Da and 65 kDa (Rosenberg and Damus, J. Biol. Chem. 248: 6490-505 (1973); Nordenman et al., Eur. J. Biochem., 78: 195-204 (1977); Kurachi et al., Biochemistry 15: 373-7 (1976)), of which some ten percent is contributed by four glucosamine-base carbohydrate chains (Kurachi et al., Biochemistry 15: 373-7 (1976); Petersen et al., in The Physiological Inhibitors of Coagulation and Fibrinolysis (Collen et al., eds) Elsevier, Amsterdam, p. 48 (1979)).

[0030] Although its name implies that it works only on thrombin, ATIII actually serves to inhibit virtually all of the coagulation enzymes to at least some extent. The primary enzymes it inhibits are factor Xa, factor IXa, and thrombin (factor IIa). It also has inhibitory actions on factor XIIa, factor Xia and the complex of factor VIIa and tissue factor but not factor VIIa and activated protein C. ATIII also inhibits trypsin, plasmin and kallikrein (Charlotte and Church, Seminars in Hematology 28:3-9 (1995). Its ability to

WO 2006/014519

PCT/US2005/024046

limit coagulation through multiple interactions makes it one of the primary natural anticoagulant proteins.

[0031] ATIII acts as a relatively inefficient inhibitor on its own. However, ATIII can be activated by a simple template mechanism, or by an allosteric conformational change brought about by heparin binding (Skinner et al., J. Mol. Biol. 283: 9-14 (1998); 5 Huntington et al., J. Mol. Biol., 293: 449-55 (1999); Belar et al., J. Mol. Biol. Chem., 275: 8733-41 (2000)). When ATIII binds heparin, the speed with which the reaction that causes inhibition occurs is greatly accelerated. This interaction is the basis of heparin based anticoagulation therapies.

10 [0032] Pure plasma derived or recombinant ATIII is inactive in lowering HIV viral loads. However, we have shown in U.S.S.N. 10/436,872 that ATIII that has been treated to have a higher molecular weight, or "activated," effectively reduces HIV viral loads in infected cells. This application, which is hereby incorporated by reference in its entirety, features pharmaceutical compositions comprising high molecular weight antithrombin III (ATIII) in 15 an amount effective to treat a retroviral infection. Incubation of the ATIII with heparin and/or heat in a variety of combinations was shown to activate the ATIII and inhibit proliferation of the retroviral infection.

[0033] The present invention is based on the surprising finding that ATIII may be activated using saccharides such as heparin directly *in situ* in a blood product, such as blood, plasma, 20 serum albumin, recombinant plasma, and the like. The blood product so treated contains activated or high molecular weight ATIII. This novel treatment eliminates the need for advance ATIII preparation, e.g. preparation, purification and formulation into a pharmaceutical composition. Such a treatment may be useful, for example, in treating patients having viral disease or diseases or conditions that are caused by or contributed to 25 by thrombin activation.

[0034] For example, the patient's own blood or plasma or serum albumin in combination with heparin may be incubated at moderate conditions (about 37 to about 40°C), (not necessary if the saccharide concentration is much lower than the ATIII equimolar concentration in the treated blood), and the activated blood product injected back into the 30 patient. The rationale behind this idea is that most patients will have between about 100 to about 150 mg ATIII per liter of blood, for which only a low heparin quantity is needed for activation.

WO 2006/014519

PCT/US2005/024046

[0035] Accordingly, a method for preparing activated ATIII *in situ* in a blood product may comprise: (a) adding to a blood product comprising ATIII a saccharide able to activate ATIII in an amount sufficient to activate said ATIII; and (b) incubating the resulting mixture of the blood product and saccharide under conditions sufficient to activate said ATIII. In certain embodiments, the saccharide may be added in a quantity in excess of that quantity of ATIII in the blood product, for example, in an about 0.1 to about 1.0 equimolar amount of the quantity of ATIII in the blood product.

[0036] The conditions sufficient to activate the ATIII are, in certain embodiments about 37°C to about 40°C for about 24 to about 72 hours. In certain embodiments, the temperature is about 37°C to about 38°C, about 38°C to about 39°C or about 39°C to about 40°C. In certain embodiments, the incubation time is about 24 to about 36 hours, about 36 to about 48 hours, about 48 to about 60 hours, or about 60 to about 72 hours.

[0037] The blood product may be any product or substance that is blood or is derived from blood. For example, blood products include, but are not limited to, whole blood, plasma, serum, serum albumin preparations, and artificial preparations of the same, such as recombinant plasma.

[0038] The saccharide, may be for example, an oligosaccharide, such as any form of heparin, including low molecular weight heparin (about 2 to about 4 kDa), high molecular weight heparin (at least 12 kDa), and standard unfractionated heparin. In certain embodiments, the saccharide is a pentasaccharide. Further, the saccharide may be an oligosaccharide treated by a glycosidase or other restriction enzyme. In certain embodiments, the saccharide may be added in an about a 0.1 to about 0.2, about 0.2 to about 0.3, about 0.3 to about 0.4, about 0.4 to about 0.5, about 0.5 to about 0.6, about 0.6 to about 0.7, about 0.7 to about 0.8, about 0.8 to about 0.9, or about 0.9 to about 1.0 equimolar amount of the quantity of ATIII in the blood product. Saccharides that may be used in the methods of the present invention include, but are not limited to, monosaccharides, disaccharides, and polysaccharides (including penta-, hepta- and hexa-saccharides), sugar alcohols, and amino sugars. Examples of monosaccharides include glucose, fructose, galactose, mannose, arabinose, and inositol. Examples of disaccharides include saccharose, lactose, maltose, pectin. Examples of sugar alcohols include mannitol, sorbitol, and xylitol. Examples of amino sugars include glucosamine, galactosamine, N-acetyl-D-glucosamine and N-acetyl galactosamine, which are the building blocks that can form more complex oligosaccharides, such as aminoglycosides and heparin. In certain embodiments, the

WO 2006/014519

PCT/US2005/024046

oligosaccharides may be low molecular weight (2-4kDa) heparin, high molecular weight (at least 12kDa) heparin, standard unfractionated heparin, pectin, pentasaccharides, and aminoglycosides. In certain embodiments, the oligosaccharide has an affinity for ATIII. Saccharides as used herein can be derivatized with additional small molecules, such as

5 biotin, avidin or streptavidin. Saccharides in certain embodiments may be sulfated oligosaccharides or oligosaccharides identified by glycosidase and other restriction enzyme reactions.

[0039] Blood products in which ATIII has been activated *in situ* are also within the scope of the present invention. Such blood products may be packaged, for example, in sterile i.v.

10 or incubation bags.

[0040] 3. Methods of Treating

[0041] The above-described methods of treating blood products to activate ATIII *in situ* and activated blood products produced thereby may be incorporated into methods of treating a disease in a subject. In certain embodiments, such methods of treating a disease

15 in a subject in need of treatment may comprise (a) adding to a blood product comprising ATIII a saccharide able to activate ATIII in an amount sufficient to activate said ATIII; (b) incubating the resulting mixture of the blood product and saccharide under conditions sufficient to activate said ATIII; and (c) administering said mixture into said subject to treat a disease. Such methods may, in certain embodiments, further comprise purifying said

20 mixture to remove unreacted saccharide prior to the infusing step, e.g. by dialysis. The methods of the present invention can be used to human and animal subjects such as cows, horses, dogs, cats, etc.

[0042] In certain embodiments, the disease is caused by a bacteria or virus. Exemplary viruses that may be treated using the methods of activating ATIII *in situ* or with the blood

25 products of the invention include Hepatitis A Virus (HAV) infection, Hepatitis B Virus (HBV) infection, Hepatitis C Virus (HCV) infection, Human Immunodeficiency Virus (HIV) infection, corona virus infection, cytomegalovirus infection (CMV) and severe acute respiratory syndrome (SARS). In certain embodiments, the virus is a retrovirus, such as HIV. In other embodiments, the disease may be a disease or condition that is caused by or

30 contributed to by thrombin activation.

[0043] Thrombin activation related diseases in a patient include sepsis, trauma, acute respiratory distress syndrome, thrombosis, stroke, and restenosis. The methods may also be used to treat patients at risk of a thrombin related pathological disease such as reocclusion

WO 2006/014519

PCT/US2005/024046

- and restenosis in percutaneous transluminal coronary angioplasty; thrombosis associated with surgery, ischemia/reperfusion injury; and coagulation abnormalities in cancer or surgical patients. The treated blood product administered in the methods may serve an anti-coagulant in the treatment of, for example, congenital antithrombin III deficiency which
- 5 leads to an increased risk of venous and arterial thrombosis, or acquired antithrombin III deficiency which results in disseminated intravascular coagulation, microangiopathic hemolytic anemias due to endothelial damage (i.e. hemolytic-uremic syndrome) and veno-occlusive disease (VOD). The methods and blood products may also be used to treat semi-chronic diseases like arterial thrombus and deep vein thrombosis.
- 10 [0044] Both methods of treating that are single-dose administrations and longer term, chronic, multi-dose treatments are within the scope of the present invention.
- [0045] In certain embodiments, the methods of treating may further comprise administering other pharmaceutical compositions, such as an anti-viral drug, to said subject. For example, in certain embodiments, an anti-viral drug may be administered concurrently with the blood
- 15 product. In other embodiments, however, an anti-viral drug may be administered subsequent to administering the mixture. In still other embodiments, anti-viral drug may be administered both concurrently and subsequently with the blood product. An anti-viral drug may be added to the mixture, or administered as a separate composition.
- [0046] Exemplary anti-viral drugs include reverse transcriptase inhibitors such as
- 20 zidovudine, zalcitabine, didanosine, stavudine, lamivudine, abacavir, tenofovir, nevirapine, efavirenz, delavirdine; protease inhibitors such as saquinavir, ritonavir, indinavir, nelfinavir, amprenavir, lopinavir, and other agents such as adenine arabinoside, adenine arabinoside 5'-monophosphate, acyclovir, ganciclovir, famciclovir, lamivudine, clevudine, afedovir dipivoxil, entecavir, IFN-a-2b, IFN-a-2a, lymphoblastoid IFN, consensus-IFN,
- 25 IFN-b, IFN-g, pegylated IFN-a-2a, corticosteroids, or thymosin a1, IL-2, IL-12, ribavirin, cyclosporin or granulocyte macrophage colony stimulating factor, fusion inhibitors such as T-20 (enfuvirtide), zinc finger inhibitors, and ribavirin.
- [0047] Highly Active Antiretroviral Therapy (HAART) is a recommended treatment for HIV and other viral infections. HAART combines two or more anti-viral medications in a
- 30 daily regimen, also known as a "cocktail." Cocktails of anti-viral drugs are well-known to those of skill in the art. For example, effective known anti-HIV cocktails include, but are not limited to, combinations such as AZT, 3TC and efavirenz; nevirapine, stavudine and lamivudine; emtricitabine and tenofovir disoproxil fumarate; indinavir, zidovudine and

WO 2006/014519

PCT/US2005/024046

lamivudine (3TC); and zidovudine and 3TC. An effective anti-HCV cocktail is ribavirin and interferon.

[0048] The combined use of blood products of the present invention and other anti-virals may reduce the required dosage for any individual component because the onset and
5 duration of effect of the different components may be complimentary. In such combined therapy, the different active agents may be delivered together or separately, and simultaneously or at different times within the day.

[0049] Other pharmaceutical compositions that may be administered either concurrently, subsequently, or both concurrently and subsequently include ATIII and interferon or
10 interferon derived drugs. For example, the blood product may be supplemented with additional ATIII prior to adding the oligosaccharide to activate it. In another embodiment, anticoagulants may be used alone or in combination with saccharides such as heparin or a pentasaccharide to treat the blood products of the invention prior to their use in a treatment method in order to improve their efficacy.

[0050] Further, high molecular weight ATIII pharmaceutical compositions as described in
15 Pending U.S. Patent Application 10/436,872 may be used to supplement the method of treatment, e.g. by administering it either concurrently, subsequently, or both concurrently and subsequently with administration of the treated blood products.

[0051] The blood products of the present invention may be administered directly to a
20 subject or may be formulated in a conventional manner using one or more physiologically acceptable carriers or excipients. For such therapy, the compounds of the invention can be formulated for a variety of loads of administration, including systemic and topical or localized administration. Techniques and formulations generally may be found in Remington's Pharmaceutical Sciences, Meade Publishing Co., Easton, PA. For systemic
25 administration, injection is preferred, including intramuscular, intravenous, intraperitoneal, and subcutaneous. The blood product may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. For injection, the blood products of the invention can be formulated in liquid solutions, preferably in physiologically compatible buffers such as Hank's solution or Ringer's solution. In addition, the blood
30 products may be formulated in solid form and redissolved or suspended immediately prior to use. Lyophilized forms of the blood products are also included.

[0052] Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The formulations may take such

WO 2006/014519

PCT/US2005/024046

forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

5 [0053] Furthermore, as those skilled in the art will understand, the dosage of any blood product, agent, compound, drug, etc. used in the methods of the present invention will vary depending on the symptoms, age and body weight of the patient, the nature and severity of the disorder to be treated or prevented, the route of administration, and the form of the supplement. Any of the subject formulations may be administered in any suitable dose,
10 such as, for example, in a single dose or in divided doses. Dosages for the blood products of the present invention, alone or together with any other compound of the present invention, or in combination with any compound deemed useful for the particular disorder, disease or condition sought to be treated, may be readily determined by techniques known to those of skill in the art, based on the present description, and as taught herein. Also, the
15 present invention provides mixtures of more than one subject compound, as well as other therapeutic agents.

[0054] ATIII has been shown to be well-tolerated when administered at a dose of ~100U/kg/day (Warren *et al.*, *JAMA* 286: 1869-78 (2001)) and has an overall elimination half-life with 18.6 h was demonstrated (Ilias *et al.* *Intensive Care Medicine* 26: 7104-7115
20 (2000)). While the dose is appropriately determined depending on symptom, body weight, sex, animal species and the like, it is generally 1-1,000 units/kg body weight/day, preferably 10-500 units/kg body weight/day of ATIII (as contained in the treated blood products of the invention) for a human adult, which is administered in one to several doses a day. In the case of intravenous administration, for example, the dose is preferably 10-100
25 units/kg body weight/day.

[0055] The precise time of administration and amount of any particular product or compound that will yield the most effective treatment in a given patient will depend upon the activity, pharmacokinetics, and bioavailability of a particular compound, physiological condition of the patient (including age, sex, disease type and stage, general physical
30 condition, responsiveness to a given dosage and type of medication), route of administration, and the like. The guidelines presented herein may be used to optimize the treatment, e.g., determining the optimum time and/or amount of administration, which will require no more than routine experimentation consisting of monitoring the subject and

WO 2006/014519

PCT/US2005/024046

adjusting the dosage and/or timing. In certain embodiments, the administration regimen may also be dependent on the relative amount of ATIII in the blood product, for example, when the blood product is the patient's blood..

5 [0056] While the subject is being treated, the health of the patient may be monitored by measuring one or more relevant indices at predetermined times during a 24-hour period. Treatment, including supplement, amounts, times of administration and formulation, may be optimized according to the results of such monitoring. The patient may be periodically reevaluated to determine the extent of improvement by measuring the same parameters, the first such reevaluation typically occurring at the end of four weeks from the onset of
10 therapy, and subsequent reevaluations occurring every four to eight weeks during therapy and then every three months thereafter. Therapy may continue for several months or even years, with a minimum of one month being a typical length of therapy for humans. Adjustments to the amount(s) of agent administered and possibly to the time of administration may be made based on these reevaluations. In embodiments wherein the
15 blood products are administered along with a HAART regimen, the blood products may be administered in the HAART rest interval.

[0057] Treatment may be initiated with smaller dosages which are less than the optimum dose of the product or compound. Thereafter, the dosage may be increased by small increments until the optimum therapeutic effect is attained.

20 [0058] 4. Kits

[0059] The present invention also provides kits comprising compositions of the present invention, and optionally instructions for their use. For example, kits for the practice of certain of the treatment methods of the invention may comprise a saccharide able to activate ATIII and instructions for use. Still other kits for the practice of certain of the treatment
25 methods of the invention may comprise blood products in which ATIII has been activated *in situ*. Further, the present invention provides kits for the methods of preparing activated ATIII *in situ* in a blood product. For example, such kits may comprising an oligosaccharide able to activate ATIII and instructions for use.

[0060] Kit components may be packaged for either manual or partially or wholly automated
30 practice of the foregoing methods. In other embodiments involving kits, this invention contemplates a kit including compositions of the present invention, and optionally instructions for their use. In other embodiments, a kit may further comprise controls,

WO 2006/014519

PCT/US2005/024046

reagents, buffers, and/or instructions for use. Such kits may have a variety of uses, including, for example, imaging, diagnosis, therapy, and other applications.

EXEMPLIFICATION

5 [0061] The invention now being generally described, it will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit the invention.

[0062] Example 1: Preparation Procedures for *In vitro* Plasma ATIII Activation Using

Standard, Unfractionated Heparin

10 [0063] Standard, unfractionated heparin for human use (1,000-10,000 units, or about 2 to about 40 mg) was added to 500 mL samples of commercially available human plasma (containing an average of about 120 to about 150 mg of ATIII) in a sterile incubation bag. The amount of heparin added to the bag depends on the viral load of the patient and the
15 treatment regimen desired. The resulting mixture was incubated at room temperature, 37°C or 40°C, for 24-72 hours. Gentle mixing was applied as necessary during the incubation period. The incubated mixture may be infused into a patient with or without further purification to remove unreacted heparin. In these procedures, the heparin dosage is very low compared to the level of ATIII in the plasma so that free heparin in the resultant
20 mixture is avoided, thus decreasing the risk for bleeding.

[0064] Example 2: Preparation Procedures for *In vitro* Blood ATIII Activation Using Standard, Unfractionated Heparin

[0065] Standard, unfractionated heparin for human use (1,000-20,000 units, or about 2 to about 40 mg) was added to 1,000 mL samples of blood drawn from HIV or HCV patients
25 (containing an average of about 70 to about 150 mg of ATIII) in a sterile incubation bag. The resulting mixture was incubated at room temperature, 37°C or 40°C, for 24-72 hours. Gentle mixing was applied as necessary during the incubation period. The incubated mixture may be infused into a patient with or without further purification to remove unreacted heparin. In these procedures, the heparin dosage is very low compared to the
30 level of ATIII in the plasma so that free heparin in the resultant mixture is avoided, thus decreasing the risk for bleeding.

[0066] Example 3: Preparation Procedures for *In vitro* Plasma ATIII Activation Using Low Molecular Weight Heparin

WO 2006/014519

PCT/US2005/024046

[0067] Low molecular weight (LMW) heparin for human use (1,000-10,000 units, or about 2 to about 20 mg) was added to 500 mL samples of commercially available human plasma (containing an average of about 70 to about 150 mg of ATIII) in a sterile incubation bag. The resulting mixture was incubated at room temperature, 37°C or 40°C, for 24-72 hours. Gentle mixing was applied as necessary during the incubation period. The incubated mixture may be infused into a patient with or without further purification to remove unreacted heparin. In these procedures, the heparin dosage is very low compared to the level of ATIII in the plasma so that free heparin in the resultant mixture is avoided, thus decreasing the risk for bleeding.

10 **[0068] Example 4: Preparation Procedures for *In vitro* Blood ATIII Activation Using Low Molecular Weight Heparin**

[0069] Low molecular weight (LMW) heparin for human use (1,000-20,000 units, or about 2 to 20 mg) was added to 1,000 mL samples of blood drawn from HIV or HCV patients (containing an average of about 70 to 150 mg of ATIII) in a sterile incubation bag. The resulting mixture was incubated at room temperature, 37°C or 40°C, for 24-72 hours. Gentle mixing was applied as necessary during the incubation period. The incubated mixture may be infused into a patient with or without further purification to remove unreacted heparin. In these procedures, the heparin dosage is very low compared to the level of ATIII in the plasma so that free heparin in the resultant mixture is avoided, thus decreasing the risk for bleeding.

20 **[0070] Example 5: Activation of ATIII in the Blood or Blood Plasma Using Heparin, Heparin Derivatives, or other Oligosaccharides Able to Activate ATIII**

[0071] ATIII may also be activated in the blood or blood plasma using the protocols described in Examples 1-4 above using a 0.1 to 1.0 equimolar amount of any heparin-derived pentasaccharide or other oligosaccharide able to activate ATIII. Examples of such are provided in the Detailed Description above.

25 **[0072] Example 6: Estimation of Protein Bound Heparin and Free Heparin in Modified Forms**

[0073] To estimate the free heparin and protein-bound heparin, both UV and refractive index (RI) integrations were used as follows.

30 [0074] To correct for the differences in the absorption detection by both UV and RI detectors, a correction coefficient was established from a pure protein standard measure with both detectors according to the following formula:

WO 2006/014519

PCT/US2005/024046

[0075] $K = UV_{\text{protein}}/RI_{\text{protein}}$ or $RI_{\text{protein}} = UV_{\text{protein}}/K$

[0076] The total RI integration may be expressed as follows:

[0077] $RI_{\text{total}} = RI_{\text{protein}} + RI_{\text{heparin}}$

[0078] Therefore, the bound heparin (B_{heparin}) is

5 [0079] $\%B_{\text{heparin}} = [(RI_{\text{total}} - UV_{\text{protein}}/K)/RI_{\text{total}}] \times 100$

[0080] Free heparin (F_{heparin}) is calculated according to:

[0081] $\%F_{\text{heparin}} = [F_{\text{heparin}}/(F_{\text{heparin}} + B_{\text{heparin}})] \times 100$

[0082] **Example 7: Assay for Inhibition of HIV-1 Replication**

[0083] X4 HTLV-IIIB (hereinafter X4 HIV; Chang *et al.*, NATURE, 363: 466-9 (1993)), a
10 prototypical T-tropic strain of HIV (American Type Tissue Collection, Monassass, VA,
USA; ATCC No. CRL-8543), was used to assess the effect of the ATIII activation
protocols on T-tropic HIV infection. The quantity of virus in a specified suspension
volume (e.g. 0.1 ml) that will infect 50% of a number (n) of cell culture microplate wells, or
tubes, is termed the Tissue Culture Infectious Dose 50 [TCID₅₀]. TCID₅₀ is used as an
15 alternative to determining virus titer by plaqueing (which gives values as PFUs or
plaque-forming units). Human T lymphoblastoid cells (H9 cells) expressing the human
leukocyte antigen proteins (HLA) B6, Bw62, and Cw3 were acutely infected with X4 HIV
at a MOI of 1×10^{-2} TCID₅₀ per milliliter. The infected H9 cells were resuspended to $5 \times$
 10^5 cells/ml in R20 cell culture medium. Two milliliters of this suspension was pipetted
20 into each well of a 24-well microtiter plate. These cells were then cultured in the presence
or absence of samples having activated ATIII for up to 12 days. Every three days (days 3,
6, 9 and 12), 1 ml cell supernatant was removed from test wells and replaced with an equal
volume of R20 cell culture medium. Control wells were similarly sampled but received
media containing samples that were untreated.

25 [0084] The concentration of the viral core protein p24 (gag) for HIV (Alliance[®] HIV-1 p24
ELISA kit, NEN[®] Life Science, Boston MA, USA) was measured for each sample obtained
at days 0, 3, 6, 9 and 12 respectively.

[0085] The results which are shown in FIGURE 1 demonstrate HIV-1 inhibitory activity in
blood treated with LMW heparin.

30

EQUIVALENTS

[0086] While specific embodiments of the subject invention have been discussed, the above
specification is illustrative and not restrictive. Many variations of the invention will

WO 2006/014519

PCT/US2005/024046

become apparent to those skilled in the art upon review of this specification and the claims below. For example, variants on the quantities of reactants given in the above Examples are within the scope of the invention, as are variants on the incubation time. The full scope of the invention should be determined by reference to the claims, along with their full scope
5 of equivalents, and the specification, along with such variations.

[0087] All publications and patents mentioned herein, including those listed below, are hereby incorporated by reference in their entirety as if each individual publication or patent was specifically and individually indicated to be incorporated by reference. In case of conflict, the present application, including any definitions herein, will control.

10 [0088] Wright, BIOASSAY, 18: 453-64 (1996); Skinner et al., J. Mol. Biol. 283: 9-14 (1998); Huntington et al., J. Mol. Biol. 293: 449-55 (1999); Deeks, *JAMA*, 286: 224-6 (2001); Stephenson, *JAMA*, 277: 614-6 (1997); Carr et al., *Lancet*, 351: 1881-3 (1998)

WO 2006/014519

PCT/US2005/024046

CLAIMS*We claim:*

1. A method of treating a disease in a subject in need of treatment for said disease,
5 comprising:
 - (a) adding to a blood product comprising ATIII a saccharide able to activate ATIII in an amount sufficient to activate said ATIII;
 - (b) incubating the resulting mixture of the blood product and saccharide under conditions sufficient to activate said ATIII; and
 - 10 (c) administering said mixture into said subject.
2. The method of claim 1, wherein said saccharide is an oligosaccharide.
3. The method of claim 2, wherein said oligosaccharide is heparin and wherein said
15 heparin is selected from the group consisting of: low molecular weight heparin, high molecular weight heparin, and standard unfractionated heparin.
4. The method of claim 2, wherein said oligosaccharide is added in a quantity in excess of that quantity of ATIII in said blood product.
- 20 5. The method of claim 1, wherein said conditions sufficient to activate said ATIII are about 37°C to about 40°C for about 24 to about 72 hours incubation.
6. The method of claim 2, wherein said oligosaccharide is added in a 0.1-1.0 equimolar
25 amount of the quantity of ATIII in said blood product.
7. The method of claim 1, wherein said blood product is blood plasma or HSA.
8. The method of claim 1, further comprising the step of purifying said mixture to
30 remove unreacted saccharide prior to said administering step.
9. The method of claim 1, wherein said disease is a viral infection selected from the group consisting of: Hepatitis A Virus (HAV) infection, Hepatitis B Virus (HBV) infection,

WO 2006/014519

PCT/US2005/024046

Hepatitis C Virus (HCV) infection, Human Immunodeficiency Virus (HIV) infection, and corona virus infection.

10. The method of claim 9, wherein said disease is Human Immunodeficiency Virus (HIV) infection.

11. The method of claim 1, wherein said disease is a disease or condition that is caused by or contributed to by thrombin activation selected from the group consisting of: sepsis, trauma, acute respiratory distress syndrome, thrombosis, stroke, restenosis, reocclusion and restenosis in percutaneous transluminal coronary angioplasty; thrombosis associated with surgery, ischemia/reperfusion injury; and coagulation abnormalities in cancer or surgical patients, an antithrombin III deficiency, venous or arterial thrombosis, disseminated intravascular coagulation, microangiopathic hemolytic anemias and veno-occlusive disease (VOD).

12. The method of claim 9, further comprising administering an anti-viral drug to said subject.

13. The method of claim 12, wherein said administering of said anti-viral drug is concurrent with said administering of said mixture.

14. The method of claim 12, wherein said administering of said anti-viral drug is subsequent to said administering of said mixture.

15. The method of claim 12, wherein an anti-viral drug is also added to said resultant mixture before administration.

16. The method of claim 1, further comprising supplementing said blood product with additional ATIII prior to adding said saccharide.

17. The method of claim 1, further comprising administering an interferon or interferon derived drug to said subject.

WO 2006/014519


PCT/US2005/024046

18. The method of claim 9, wherein said administering of said mixture results in a reduction of viral load.
19. A method for preparing a blood product comprising activated ATIII, comprising:
- 5 (a) adding to a blood product comprising ATIII a saccharide able to activate ATIII in an amount sufficient to activate said ATIII; and
- (b) incubating the resulting mixture of the blood product and saccharide under conditions sufficient to activate said ATIII.
- 10 20. A kit for preparing activated ATIII *in situ* in a blood product comprising an oligosaccharide able to activate ATIII and instructions for use.

WO 2006/014519

1/2

PCT/US2005/024046
FIGURE 1




Heparin activity is due to BSA in the incubation sample. This activity decreases after day nine and is zero on day 12.

WO 2006/014519

2/2

PCT/US2005/024046



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US05/24046

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : C12Q 1/00 US CL : 435/4 According to International Patent Classification (IPC) or to both national classification and IPC												
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/4 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Please See Continuation Sheet												
C. DOCUMENTS CONSIDERED TO BE RELEVANT												
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.										
X	BAYER CORPORATION, Thrombate III, Antithrombin III (Human) data sheet, last revised. 2001, pages 1-7, see entire document.	1-20										
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.												
* Special categories of cited documents: <table border="0"> <tr> <td>"A" document defining the general state of the art which is not considered to be of particular relevance</td> <td>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>"E" earlier application or patent published on or after the international filing date</td> <td>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>"O" document referring to an oral disclosure, use, exhibition or other means</td> <td>"&" document member of the same patent family</td> </tr> <tr> <td>"P" document published prior to the international filing date but later than the priority date claimed</td> <td></td> </tr> </table>			"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family	"P" document published prior to the international filing date but later than the priority date claimed	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention											
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone											
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art											
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family											
"P" document published prior to the international filing date but later than the priority date claimed												
Date of the actual completion of the international search 07 December 2005 (07.12.2005)		Date of mailing of the international search report 19 DEC 2005										
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US Commissioner for Patents P.O. Box 1450 Alexandria, Virginia 22313-1450 Facsimile No. (571) 273-3201		Authorized officer <i>Stacy B. Chem</i> Telephone No. 571.272.1600										

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US05/24046

Continuation of B. FIELDS SEARCHED Item 3:
EAST, MEDLINE, BIOSIS, EMBASE; CAPLUS; SCISEARCH
search terms: elmahleh, atIII, anti-thrombin, saccharide, hiv, blood, hav; hbv